

IN THE CLAIMS:

Kindly rewrite Claims 1-31 as follows, in accordance with 37 C.F.R. § 1.121:

1. (currently amended) An isolated strain of *Methylophilus methylotrophus* having L-lysine-producing ability, wherein dihydrodipicolinate synthase activity is enhanced as compared to a wild-type *Methylophilus methylotrophus* strain, and wherein said dihydrodipicolinate synthase is selected from the group consisting of:

a) a protein encoded by the DNA sequence depicted in a nucleotide sequence comprising nucleotide numbers 1268 to 2155 of SEQ ID NO:9; and

b) a protein having dihydrodipicolinate synthase activity and encoded by a DNA sequence which hybridizes to the DNA sequence depicted in a nucleotide sequence comprising nucleotide numbers 1268 to 2155 of SEQ ID NO:9 under stringent conditions, wherein said conditions comprise washing at 60°C in a salt solution of 1xSSC and 0.1%SDS.

2. (currently amended) The isolated strain according to claim 1, wherein the L-amino acid is L-lysine.

3. (cancelled)

4. (cancelled)

5. (currently amended) An isolated strain of *Methylophilus methylotrophus* having L-lysine-producing ability, wherein dihydrodipicolinate synthase activity and aspartokinase activity are enhanced as compared to a wild-type *Methylophilus methylotrophus* strain, and wherein said dihydrodipicolinate synthase is selected from the group consisting of:

a) a protein encoded by the DNA sequence depicted in a nucleotide sequence comprising nucleotide numbers 1268 to 2155 of SEQ ID NO:9; and

b) a protein having dihydrodipicolinate synthase activity and encoded by a DNA sequence which hybridizes to the DNA sequence depicted in a nucleotide sequence

comprising nucleotide numbers 1268 to 2155 of SEQ ID NO:9 under stringent conditions, wherein said conditions comprise washing at 60°C in a salt solution of 1xSSC and 0.1%SDS,

and wherein said aspartokinase is selected from the group consisting of:

a) a protein encoded by the DNA sequence depicted in a nucleotide sequence comprising a-nucleotide numbers 510 to 1736 of SEQ ID NO:5;

b) a protein having aspartokinase activity and encoded by a DNA sequence which hybridizes to the DNA sequence depicted in a nucleotide sequence comprising nucleotide numbers 510 to 1736 of SEQ ID NO:5 under stringent conditions, wherein said conditions comprise washing at 60°C in a salt solution of 1xSSC and 0.1%SDS.

6. (cancelled)

7. (currently amended) An isolated strain of *Methylophilus methylotrophus* having L-amino acid-producing ability, wherein aspartokinase activity is enhanced as compared to wild-type *Methylophilus methylotrophus* strain, and wherein said aspartokinase is selected from the group consisting of:

a) a protein encoded by the DNA sequence depicted in a nucleotide sequence comprising a-nucleotide numbers 510 to 1736 of SEQ ID NO:5; and

b) a protein having aspartokinase activity and encoded by a DNA sequence which hybridizes to the DNA sequence depicted in a nucleotide sequence comprising nucleotide numbers 510 to 1736 of SEQ ID NO:5 under stringent conditions, wherein said conditions comprise washing at 60°C in a salt solution of 1xSSC and 0.1%SDS.

8. (previously presented) The isolated strain according to claim 5, wherein an activity or activities of one, two, or three of enzymes selected from the group consisting of aspartic acid semialdehyde dehydrogenase, dihydrodipicolinate reductase and diaminopimelate decarboxylase is/are enhanced as compared to a wild-type *Methylophilus methylotrophus* strain.

9. (currently amended) The isolated strain according to claim 5, wherein the

dihydrodipicolinate synthase activity and the aspartokinase activity are enhanced as compared to a wild-type *Methylophilus methylotrophus* strain by ~~introduction~~ introducing into cells of said DNA sequence coding for said dihydrodipicolinate synthase ~~that does not suffer from feedback inhibition by L-lysine~~ and said DNA sequence coding for said aspartokinase ~~that does not suffer from feedback inhibition by L-lysine~~.

10. (previously presented) The isolated strain according to claim 7, wherein activities of homoserine dehydrogenase, homoserine kinase and threonine synthase are enhanced as compared to wild-type *Methylophilus methylotrophus* strain, and wherein said isolated strain has L-threonine-producing ability.

11. (cancelled)

12. (currently amended) A method for producing ~~an~~ L-lysine, which comprises culturing said strain as defined in claim 1 in a medium, accumulating said L-lysine in said medium, and collecting the L-lysine from said medium.

13. (original) The method according to claim 12, wherein the medium contains methanol as a main carbon source.

14. (withdrawn)

15. (withdrawn)

16. (currently amended) ~~A~~ An isolated DNA which codes for a protein defined in the following (A) or (B) selected from the group consisting of:

(A) a protein ~~which has~~ comprising the amino acid sequence of SEQ ID NO: 6, ~~or and~~

(B) a protein ~~which has~~ comprising an amino acid sequences sequence of SEQ ID NO:6 ~~including except that~~ substitution, deletion, insertion, addition, or inversion of one or several amino acids in said amino acid sequence is present, and wherein said

protein has aspartokinase activity.

17. (currently amended) The DNA according to claim 16, ~~which is a~~wherein said DNA defined in the following (a) or (b) is selected from a group consisting of:

(a) a DNA ~~which has comprising~~ a nucleotide sequence ~~comprising the nucleotide sequence of the nucleotide numbers 510 to 1736 of SEQ ID NO:5; or and~~

(b) a DNA which is ~~hybridizable~~ able to hybridize with a probe having the nucleotide sequence of the nucleotide numbers 510 to 1736 of SEQ ID NO:5 or a part thereof under a stringent ~~condition~~conditions, and wherein said conditions comprise washing at 60°C in a salt solution of 1XSSC and 0.1%SDS, and wherein said DNA codes for a protein having aspartokinase activity.

18. (withdrawn)

19. (withdrawn)

20. (currently amended) ~~The~~ An isolated DNA which codes for a protein ~~defined in the following (E) or (F)~~selected from the group consisting of:

(E) a protein ~~which has comprising~~ the amino acid sequence of SEQ ID NO:10, ~~or and~~

(F) a protein ~~which has comprising~~ an amino acid ~~sequences~~ sequence of SEQ ID NO:10 ~~including except that~~ substitution, deletion, insertion, addition, or inversion of one or several amino acids in said amino acid sequence is present, and wherein said protein has dihydrodipicolinate synthase activity.

21. (currently amended) The DNA according to claim 20, ~~which is a~~wherein said DNA defined in the following (e) or (f) is selected from the group consisting of:

(e) a DNA ~~which has comprising~~ a nucleotide sequence ~~comprising the nucleotide sequence of the nucleotide numbers 1268 to 2155 of SEQ ID NO:9; or and~~

(f) a DNA which is ~~hybridizable~~ able to hybridize with a probe having the nucleotide sequence of the nucleotide numbers 1268 to 2155 of SEQ ID NO:9 or a part

thereof under ~~a stringent condition~~ conditions, and wherein said conditions comprise washing at 60°C in a salt solution of 1XSSC and 0.1%SDS, and wherein said DNA codes for a protein having dihydrodipicolinate synthase activity.

22. (withdrawn) The DNA which codes for a protein defined in the following (G) or (H):

(G) a protein which has the amino acid sequence of SEQ ID NO:12, or

(H) a protein which has an amino acid sequences of SEQ ID NO:12 including substitution, deletion, insertion, addition, or inversion of one or several amino acids, and has dihydrodipicolinate reductase activity.

23. (withdrawn) The DNA according to claim 22, which is a DNA defined in the following (g) or (h):

(g) a DNA which has a nucleotide sequence comprising the nucleotide sequence of the nucleotide numbers 2080 to 2883 of SEQ ID NO:11 or

(h) a DNA which is hybridizable with a probe having the nucleotide sequence of the nucleotide numbers 2080 to 2883 of SEQ ID NO:11 or a part thereof under a stringent condition, and codes for a protein having dihydrodipicolinate reductase activity.

24. (withdrawn) The DNA which codes for a protein defined in the following (I) or (J):

(I) a protein which has the amino acid sequence of SEQ ID NO:14, or

(J) a protein which has an amino acid sequences of SEQ ID NO:14 including substitution, deletion, insertion, addition, or inversion of one or several amino acids, and has diaminopimelate decarboxylase activity.

25. (withdrawn) The DNA according to claim 24, which is a DNA defined in the following (i) or (j):

(i) a DNA which has a nucleotide sequence comprising the nucleotide sequence of the nucleotide numbers 751 to 1995 of SEQ ID NO:13; or

(j) a DNA which is hybridizable with a probe having the nucleotide sequence of the

nucleotide numbers 751 to 1995 of SEQ ID NO:13 or a part thereof under a stringent condition, and codes for a protein having diaminopimelate decarboxylase activity.

26. (currently amended) The isolated strain according to claim 1, wherein an activity or activities of one, two, or three of enzymes selected from the group consisting of aspartic acid semialdehyde dehydrogenase, ~~dihydrokippiculate~~ dihydrodipicolinate reductase and diaminopimelate decarboxylase is/are enhanced as compared to a wild-type *Methylophilus methylotrophus* strain.

27. (previously presented) The isolated strain according to claim 2, wherein an activity or activities of one, two, or three of enzymes selected from the group consisting of aspartic acid semialdehyde dehydrogenase, dihydrodipicolinate reductase and diaminopimelate decarboxylase is/are enhanced as compared to a wild-type *Methylophilus methylotrophus* strain.

28. (currently amended) A method for producing ~~an~~ L-lysine, which comprises culturing said strain as defined in claim 5 in a medium, accumulating said L-lysine in said medium, and collecting the L-lysine from said medium.

29. (previously presented) The method according to claim 28, wherein the medium contains methanol as a main carbon source.

30. (previously presented) A method for producing an L-amino acid, which comprises culturing said strain as defined in claim 7 in a medium, accumulating said L-amino acid in said medium, and collecting the L-amino acid from said medium.

31. (previously presented) The method according to claim 30, wherein the medium contains methanol as a main carbon source.